

Weaver-Dunn Protocol

Surgery is generally indicated for chronic grade III and IV AC separations. The procedure is performed for functional rather than cosmetic purposes.

Recommendations:

- Elbow must be supported at all times for 6 weeks post-op. Use sling but not swathe as it may pull the arm inferiorly.
- No driving until 6 weeks post-op.
- Encourage PROM at home <u>daily</u> (2 3 sessions) by family member for the first 4 6 weeks.
- Instruct in proper posture and use of pillows to support arm while sleeping.
- Have patient ice shoulder 3 4 times daily to decrease pain and inflammation.
- <u>PROM Limits</u>: Forward elevation and abduction to 90° for 6 weeks. Internal and external rotation to tolerance.
- Return to work and sport to be determined on an individual basis by the physician.

Post-Op Protocol:

0 - 2 Weeks:

- 1. PROM in all planes. FE and Abduction limited to 90°. ER/IR to tolerance.
- 2. Begin multi-angle isometrics (submax).
- 3. AROM of elbow, wrist and hand with arm supported.

2 - 4 Weeks:

- 1. Continue progressing PROM within limitations
- 2. Scapular setting in sling.
- 3. Supine ER with wand.

4 - 6 Weeks:

1. Continue progressing PROM. Should have attained 90° FE and abduction and 65-70° IR/ER by 4 weeks.

Developed in conjunction with the physicians at OrthoCarolina

Weaver-Dunn.1 Page 1 of 2 Revised: May 2005

- 2. Begin active side lying ER.
- 3. <u>Gentle</u> open kinetic chain rhythmic stabilization progression in supine.

6 - 8 Weeks:

- 1. PROM to tolerance.
- 2. Begin AAROM with pulleys and wand.
- 3. UBE avoiding excessive protraction/retraction.
- 4. AROM FE and abduction to 90°.
- 5. Begin scapular stabilization and rotator cuff exercises within pain-free range.

8 - 12 Weeks:

- 1. Full AROM FE 170, ER 80-90, IR 90 by 12 weeks.
- 2. Progress parascapular and rotator cuff exercises.
- 3. Train extrinsic shoulder musculature (deltoids, biceps, triceps)
- 4. Plyoback, advanced PNF with theraband, bodyblade etc.

*Full unrestricted activity by 6 months.